

The effectiveness of wet cupping vs. venesection on arterial O₂ saturation level of cigarette smokers: A randomized controlled clinical trial

Hekmatpou D¹, Moeini L², Haji-Nadali S³

ABSTRACT

Objective: Wet cupping is a traditional bloodletting method recommended for controlling of respiratory disease complications. This study aimed to compare the efficacy of wet cupping vs. venesection on arterial O₂ saturation level of smokers.

Methods: This is a randomized controlled clinical trial which started with simple sampling of smokers. After administering spirometry, participants (N = 110 male smokers) with positive pulmonary function test (PFT), who manifested Chronic Obstructive Pulmonary Disease (COPD), were randomly assigned to intervention and control groups. The two groups were assessed in terms of demographic data, rate of hemoglobin (Hb), hematocrit (Hct), and arterial O₂ saturation. Then, the intervention participants underwent wet cupping whereas venesection was performed on the control participants. At four stages after the two treatments, pulse oximetry was performed. Data was analyzed using SPSS (Version 17).

Results: Result shows that mean arterial O₂ sat level increased at three stages, namely before, immediately after, and 6 and 12 hrs after these two treatments ($p \leq 0.001$). This indicates that wet cupping and venesection alike were effective on O₂ sat level in the two groups, but the increasing pattern was maintained 12 hrs afterward only in those participants who had received wet cupping ($p \leq 0.001$). Moreover, the results of repeated measure ANOVA between the two groups at the four stages showed that there were significant differences between the means of O₂ saturation level at the 6- and 12-hrs stages ($F = 66.92$, $p \leq 0.001$).

Conclusion: Wet cupping caused a continued O₂ saturation in the intervention group even up to 12 hrs afterward. Participants expressed liveliness and improved respiration after wet cupping. Therefore, wet cupping is recommended for promoting the health of cigarette smokers.

KEY WORDS: Arterial O₂ saturation, Cigarette smokers, Venesection, Wet cupping.

doi: <http://dx.doi.org/10.12669/pjms.296.3365>

How to cite this:

Hekmatpou D, Moeini L, Haji-Nadali S. The effectiveness of wet cupping vs. venesection on arterial O₂ saturation level of cigarette smokers: A randomized controlled clinical trial. *Pak J Med Sci* 2013;29(6):1349-1353.

doi: <http://dx.doi.org/10.12669/pjms.296.3365>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Hekmatpou D, PhD (Nursing), Assistant Professor,
2. Dr. Moeini L, MD, Specialist in Pulmonary Diseases, Assistant Professor,
3. Salar Haji-Nadali, Student of Nursing, Student Research Center Committee,
- 1-3: Faculty Member of Arak University of Medical Sciences, Arak, Iran.

Correspondence:

Hekmatpou D, PhD in Nursing,
E-mail: dr_hekmat@arakmu.ac.ir

- * Received for Publication: January 7, 2013
- * First Revision Received: August 12, 2013
- * Second Revision Received: August 19, 2013
- * Final Revision Accepted: August 20, 2013

INTRODUCTION

Cigarette smoking is one of the major problems of public health and premature deaths.^{1, 2} Studies have shown that smoking is becoming increasingly prevalent in Asian countries compared to the rest of the world. Studies in central Iran reported that up to 15% of Iranians over age 19 smoked.³

About 90% of patients with a COPD are smokers. One of the effects of smoking is the increasing of hemoglobin, hematocrit, and red blood cell counts, which occur as a result of changes in blood gases.⁴

Despite great advances in medical sciences and overwhelming popularity of conventional medicine, the growth and development of public interest in and use of complementary or alternative medicine (CAM) are well documented.⁵

Hijama means cupping, but in Arab and Muslim culture it refers to wet cupping.⁶ Wet cupping is over 5000 years old and it was possibly the Egyptians who first practiced bloodletting by scarification.⁷ The first precursors of cupping were Hippocrates and Galen, and after the advent of Islam, it was fully presented to humanity by Prophet Muhammad (PBUH).⁸ The mechanism of cupping therapy is not clear, but some researchers suggest that placement of cups on selected acupoints on the skin produces hyperemia or hemostasis, which results in a therapeutic effect.^{9,10}

Services and treatments provided by conventional medicine have proved costly for rehabilitated smokers. Besides, they are not totally satisfied with their treatment. These reasons and the gap in alternative medicine motivated the researchers of the present study to investigate the effects of wet cupping on arterial blood O2 saturation of smokers.

METHODS

This study is a randomized controlled clinical trial, which is registered on IRICT.ir by 201108134519N2. Smokers visiting a lung clinic in Mahalat, central Iran, were chosen by simple method of sampling (sample size power = 80%, alpha = 0.05, size effect = 50% Altman nomogram for former and future studies = 49 patients). With a 5% probability of attrition, 55 subjects for each control and intervention groups were determined. The

inclusion criteria were (a) granting formal consent, (b) not having acute respiratory illnesses or cold, (c) not having smoked for 3 - 6 hrs before, (d) and not having neuromuscular (e.g., myasthenia gravis) and metabolic (e.g. diabetes and blood) diseases. The exclusion criteria were (a) declaring disinclination halfway, (b) catching a cold, (c) smoking cigarettes during the study, (d) suffering from metabolic and hematologic diseases. After cigarette smokers were selected and agreed to participate voluntarily with written consent, they took a spirometry test.

Participants with a normal pulmonary function test were omitted and those with obstructive PFT (American Thoracic Society's index of diagnosis) were randomly divided into two intervention and control groups. Then, the two groups were compared in terms of hemoglobin, hematocrit, and arterial O2 saturation using pulse oximetry (type Critical Care 502). Afshinjoo et al showed that there was a strong correlation between the percentage of arterial O2 saturation from pulse oximetry and the percentage of O2 saturation in arterial blood test (ABG) (r = 95%).¹¹ Subsequently, wet cupping and venesection were performed on participants. Finally, 6 and 12 hrs after the intervention, pulse oximetry tests was performed. Data were analyzed using SPSS (Version 17).

In this study, wet cupping was carried out based on the common procedure. After a careful examination by a physician, if wet cupping is considered necessary and beneficial for him, the patient sits cross-legged on the bed. Then, the physician antiseptices a certain site using Betadine or alcohol and places a specially designed cup on it. After that, a suction device is used to vacuum the

Table-I: Distribution of smokers based on demographic factors.

Variable	Type	Group		Statistical test X2/ P- value
		Wet Cupping (No/%)	Venesection (No/%)	
Marriage Status	Married	46/83.6	40/72.7	1.91/0.16
	Single	6/16.4	15/27.3	
Education Level	Low Literates (Under Diploma)	38/69.1	37/67.3	8.5/ 0.075
	Diploma	14/25.5	8/14.5	
	University level	3/5.5	10/18.2	
Income*	Low	9/16.4	12/21.8	3.69/ 0.15
	Intermediate	33/60	23/41.8	
	Good	13/23.6	20/36.4	
Job	Worker	46/83.6	38/69.1	9.42/ 0.71
	Clerk	9/16.4	10/18.2	
	Retired	0/0	7/12.7	

*(Low=Income ≤ 250\$/month, Intermediate= Income = 250- 500\$/month, Good= Income ≥ 500 \$/month)

Table-II: Distribution of mean and standard deviation of special data of two groups.

Group Variable	Wet cupping Mean \pm SD	Venesection Mean \pm SD	(p-value)
Age*	50.34 \pm 6.7	50.45 \pm 5.7	0.61
Weight (kg) **	80.8 \pm 10.21	79.6 \pm 9.6	0.26
Height (cm) **	172 \pm 6.6	173 \pm 7	0.28
Body Mass Index**	27.5 \pm 3.7	26 \pm 4	0.71
Smoking duration **	14 \pm 7.6	13 \pm 6	0.65
Smoking no /day**	22 \pm 5	23 \pm 5	0.75
Diagnosis of Disease**	4 \pm 2.6 (year)	3.6 \pm 2.5(year)	0.08

* t test, ** Man Whitney U test

cup. The vacuum condition and the environmental air pressure pull the patient's skin into the cup so much so that it is firmly stuck to the skin. (The proper height of the skin dome from its top to the edge of the suction cup must be 1 to 1.5 cm.). This suction creates local congestion and inflammation, causing an accumulation of blood in the capillaries. After 3 to 5 minutes, the practitioner removes the cup and, utilizing a disposable sterile surgical razor, makes several scarifications, each 0.5 to 1 mm deep. The cup is put back on the same location and after re-suction; blood is gradually drawn out of the cuts. The cycle is repeated 3 to 5 times, each lasting for 3 to 5 min. In general, a wet cupping session takes 15 to 20 minutes. and 50 to 75 ml. blood in total is drawn out. In the control group, venesection was done using the sterile technique. The blood taken was 100 to 200 ml. The results of pulse oximetry at 6- and 12-hr stages after were compared in both groups.

Ethical considerations: The study was a research project approved in Arak University of Medical Sciences, which was also confirmed by Clinical Research Ethics Committee of the University.

RESULTS

The cohort of 110 smokers was randomly assigned to two groups, namely Wet Cupping (Hijama) and Venesection.

Results of the statistical test of X^2 did not show a significant difference between the two groups in terms of the above variables. The smoking variable was equal in both groups and all participants in the study (100%) had been smoking more than 10 years.

The mean age of both groups was about 55 years. There were no significant differences in the distribution of participants in the two groups based on the variables mentioned.

Results from the Mann-Whitney U test do not display any significant difference between the two groups before intervention. In other words, the distribution of participants in the two groups did not have any significant difference in terms of variables mentioned.

Results show that in both groups, arterial O2 saturation rose at all four stages that is before, immediately after, six hrs after and 12 hrs after wet cupping and venesection. That is to say both venesection and wet cupping were effective on O2 sat. Results of a paired t-test showed that the mean arterial O2 saturation before, after, six and twelve hrs after wet cupping were significantly different ($p \geq 0.001$); that is, the arterial O2 saturation at the four stages was on the rise. Although the results of the paired t-test of the venesection group showed that the mean arterial O2 saturation at all the four stages were significantly different ($0.01/0 \geq P$), this

Table-III: Distribution of mean and standard deviation of other special data of two groups.

Group Variable	Wet Cupping Mean \pm SD	Venesection Mean \pm SD	Man Whitney U (p-value)
Hemoglobin	17.7 \pm 0.73	17.5 \pm 0.63	0.61
Hematocrit	53 \pm 1.7	54 \pm 1.6	0.26
FVC* rate	3.9 \pm 0.53	3.5 \pm 0.81	0.28
VC* rate	3.9 \pm 0.62	3.18 \pm 0.76	0.71
FEV1/FVC* rate	0.77 \pm 0.17	0.79 \pm 0.1	0.65
FVC%	0.81 \pm 0.17	0.79 \pm 0.15	0.75
VC%	3.5 \pm 3.5	3 \pm 4.1	0.08
FEV1/FVC%	0.78 \pm 0.17	0.79 \pm 0.16	0.35

FVC* rate = Force Vital Capacity

VC* rate = Vital Capacity

FEV1/FVC* rate = Force Expiratory Volume in the first second/ Force Vital Capacity

Table-IV: Distribution of mean and standard deviation of arterial O2 Saturation of two groups.

Group Variable	Wet Cupping Mean \pm SD	Venesection Mean \pm SD	t test (p-value)
O2Sat. Before	88.5 \pm 3.7	88.5 \pm 4.1	0.75
O2Sat. Immediately after	89 \pm 3.6	89 \pm 4.2	0.88
O2Sat. 6 hours after	93.5 \pm 3	92.5 \pm 3	\leq 0.001
O2Sat. 12 hours after	94 \pm 3.3	92 \pm 3.4	\leq 0.001




Fig.1: Differences among O2 Sat.Rate in two groups.

increasing trend stopped after the 12-hr stage. There was no significant difference between the mean arterial O2 saturation levels six and twelve hrs after venesection ($p \geq 0.035$).

Doing repeated measure ANOVA test between the two groups at four stages (Wilks' Lambda), indicates a significant difference between the mean arterial O2 saturation between the two groups 6 and 12 hrs after the two measures ($p \geq 0.001$ and $F = 66.92$).

DISCUSSION

Based on wet cupping hypothesis, wet cupping bloodletting is different from venous bloodletting in that the blood is drawn from capillary tubes and includes some lymph fluid, modifying its concentration and eliminating waste materials.^{8,9} Between the two shoulders, lymphatic and venous capillary systems are built such that toxins of blood, heavy particles, and harmful wandering macro-elements are caught in the blood vessels. Wet cupping in this area of the body sucks out contaminated blood.⁸

In this study the results indicated that in both groups arterial O2 saturation at the four stages were on the rise. This means that both interventions, i.e. wet cupping and venesection, were effective on O2 saturation. The means of arterial O2 saturation at before, after, six, and twelve hrs stages of wet cupping were significantly different ($p \geq 0.0001$), meaning that the saturation of arterial O2 at the four stages was on the rise. As for the Venesection

group, there were significant differences between the means of arterial O2 saturation at the four stages ($p \geq 0.001$). However, this growing trend stopped 12 hrs afterward, and there was no significant difference between the mean arterial blood O2 saturation at 6- and 12- hr stages ($p \geq 0.035$). Apparently, this difference, especially after the 12-hr stage of wet cupping, is due to the difference in the biochemical composition of blood. A study comparing the venous and wet cupping bloods showed that the amount of LDL and TG and the levels of RBC, Hb, hematocrit, and blood viscosity were higher in the wet cupping blood.^{12, 13}

In another study, it was reported that wet cupping lead to a significant increase of HDL levels and reduction of LDL and triglycerides.^{14,15} Thomas and Wilson in their study have stated that Bernard et al demonstrated the effect of wet cupping on O2 uptake increase in patients with polycythemia.¹⁵ A comparison of venous and wet cupping blood showed that the concentration of uric acid, urea, triglycerides, and cholesterol was significantly higher in the wet cupping blood.¹⁶ Shekarforoush and Foadoddini in their study have stated that the rate of ischemic induced arrhythmias was significantly modified by wet cupping ($P < 0.05$). These results indicate for the first time in rats that cupping might be cardioprotective in the ischemic reperfusion injury model.¹⁷

ANOVA test results between the Wet cupping and Venesection Groups at the four stages (Wilks' Lambda), indicated a significant difference between the mean arterial O2 saturation in both groups at the 6- and 12-hr stages ($p \geq 0.001$ and $F = 66.92$). This discrepancy can be due to viscosity reduction following the wet cupping. Because comparatively more amounts of lipids, uric acid, and other waste material are removed from blood after wet cupping, arterial O2 saturation would be higher after wet cupping. In this regard, several studies have been conducted which show different blood density in wet cupping and venesection bloods.

In Fazel et al. study the results showed that LDL levels after wet cupping decreased significantly ($p > 0.001$). Changes in the LDL / HDL ratio using the t-test showed that this ratio had declined more

in the wet cupping group than the control group, and that there was a significant difference between the two groups ($p > 0.004$). Hence, by reducing LDL and the LDL / HDL ratio, wet cupping can play a pivotal role in eliminating the risk factors of atherosclerosis and cardiovascular diseases.¹⁶ Daniali et al in their study have stated that the amount of uric acid, HDL, TG, LDL, SGOT and iron levels were higher in the wet cupping blood, and this difference vs. venous bloods was statistically significant ($p < 0.001$). Results also showed that the amount of red blood cells, hemoglobin, hematocrit, viscosity, MCHD (Mean Corpuscular Hemoglobin Concentration) in the wet cupping blood compared with venous blood were significantly higher ($p < 0.013$).¹⁸ In one study the results suggested significant effect in favor of cupping on vascular compliance and degree of vascular filling.¹⁹

These results show that venous blood and wet cupping blood are different in terms of biochemical properties. This can be due to the area on which wet cupping and phlebotomy (venesection) were done and differences that can be associated with the adverse site or manner of wet cupping. It seems that these differences could increase the arterial O₂ saturation rate among smokers, but for having better advantages doing clinical trials with more samples is recommended.

CONCLUSION

This study provided evidence that performing wet cupping on cigarette smokers leads to enhanced arterial O₂ saturation with an increasing trend 12 hours afterward. Those participants who had received the wet cupping treatment expressed satisfaction on better breathing. Therefore, wet cupping is recommended for promoting the health of cigarette smokers.

ACKNOWLEDGMENT

The authors would like to extend their thanks to the Research Council, the University Ethics Committee, Vice President of Education and Research, Student Research Committee members, and all the students who did a favor in this research project.

REFERENCES

1. Wald PG, Vos T, Narman R, Laubscher R, Van Walbeek C, Saloojee Y, et al. Estimating the burden of disease attributable to Smoking in South Africa in 2000. *S Afr Med J*. 2007;97:674-681.
2. Lopez MJ, Perez-Rios M, Ffino AS, Fortmann SP, Jamrozik K. Mortality attributable to passive smoking in Spain, 2002. *J Tob Contr*. 2007;16:373-377.
3. Sarafzadegan N, Sadri G, Malek-Afzali H. Isfahan Healthy heart program. A comprehensive integrated community based program for cardiovascular disease. *Acta Cardiol*. 2003;58(4):304-320.
4. Rannard SI. Treatment of stable Chronic Obstructive Pulmonary Disease. *Lancet*. 2004;364(9436):791-803.
5. Elolemy AT, AlBedah A. Public knowledge, attitude and practice of complementary and alternative medicine in Riyadh region, Saudi Arabia. *Oman Med J*. 2012;27(1):20-26.
6. Albedah A, Khalil M, Elolemy A, Elsubai I, Khalil A. Hijama (cupping): A review of the evidence (Review). *Foc Alt Compl Ther*. 2011;16(1):12-16.
7. Parapia LA. History of bloodletting by phlebotomy. *Br J Haematol*. 2008;143(4):490-495.
8. Gharebaghian A, Mehran M, Karimi GH, Vafaeiyan V, Namini MT. History of Hijama. *Blood Quar*. 2008;6(2):147-158. (Persian).
9. Cao H, Li X, Liu J. An Updated Review of the Efficacy of Cupping Therapy. *Plos One*. 2012;7(2):1-14.
10. Cao H, Han M, Li X, Dong S, Shang Y, Wang Q, Xu S, Liu L. Clinical research evidence of cupping therapy in China: a systematic literature review. *BMC Compl Alt Med*. 2010;10(70):1-10.
11. Afshin Jo M, Hamzeh Pour MR, Taghilo Gh. Comparing of two pulse oximetry and arterial blood gases tests for determining of arterial O₂ Saturation. *ZJMS*. 2002;44:25-29. (Persian).
12. Agha Rafiee AA. The Center for Islamic Medicine, Tehran, Iran. 2nd International Congress on Traditional Medicine and Materia Medica. *I J Pharm Res*. 2004;66-78. (Persian).
13. Niasari M, Kosari F, Ahmadi A. The effect of wet cupping on serum lipid concentrations of clinically healthy young men: A randomized controlled trial. *J Alt Compl Med*. 2007;13(1):79-82.
14. Montazeri R. Comparing of biomedical compositions of venous blood with Hijama blood. PhD. Dissertation no: 7522. Shahid Beheshti University. 2002. (Persian).
15. Thomas DJ, Wilson JR. The Effect of phlebotomy on renal function and proteinuria in a patient with congenital cyanotic heart disease. *Nephron*. 1983;33(3):225-226.
16. Fazel A, Hossani Vaez Z, Saghebi A, Esmaeili H. Effect of Wet cupping on serum lipoproteins concentrations of patients with hypercholesterolemia. *Mashhad J Nur Midwif*. 2008;9(1):13-18. (Persian).
17. Shekarforoush S and Foadoddini M. Cardiac effects of cupping: myocardial infarction, arrhythmias, heart rate and mean arterial blood pressure in the rat heart. *Chin J Physiol*. 2012;55(4):253-258.
18. Daniali F, Vaeze Mahdavi MR, Ghazanfari T, Naseri M. Comparing of venous blood with Hijama blood according to biochemical and hematological factors and immunological responses. *J Phy Phar*. 2008;13(1):78-87. (Persian).
19. Lee MS, Choi TY, Shin BC, Kim JI, Nam SS. Cupping for hypertension: a systematic review. *Clin Exp Hypertens*. 2010;32(7):423-425.

Author Contribution:

DH, LM and SHN: Conceived, designed and did statistical analysis & editing of manuscript.

DH, LM and SHN: Did data collection and manuscript writing.

DH, LM and SHN: Did review and final approval of manuscript.